

(3 hours)

Total marks: 80

N.B.: (1) Question No. 1 is compulsory

(2) Attempt any Three from remaining

Q1 a) If $\log \tan x = y$ then prove that $\sinh ny = \frac{1}{2} [\tan^n x - \cot^n x]$ [3]b) If $u = x^2y + e^{xy^2}$ Find $\frac{\partial^2 u}{\partial x \partial y}$ [3]c) If $x = u - uv$, $y = uv - uvw$, $z = uvw$ find $\frac{\partial(x,y,z)}{\partial(u,v,w)}$ [3]d) Using MacLaurin's series, Prove $e^{ex} = e + ex + ex^2 + \dots$ [3]e) Show that $A = \begin{bmatrix} \alpha + i\gamma & -\beta + i\delta \\ \beta + i\delta & \alpha - i\gamma \end{bmatrix}$ is unitary if $\alpha^2 + \beta^2 + \gamma^2 + \delta^2 = 1$ [4]f) Find n^{th} derivative of $\frac{x}{(x-1)(x-2)(x-3)}$ [4]Q2 a) Solve $x^5 = 1 + i$ and find the continued product of the roots. [6]b) Reduce the matrix $A = \begin{bmatrix} 5 & -6 & -6 \\ -1 & 4 & 2 \\ 3 & -6 & -4 \end{bmatrix}$ to the normal form and find its Rank [6]

c) State and Prove Euler's theorem for two variables hence [8]

find value of $x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y}$ where $u = \frac{\sqrt{xy}}{\sqrt{x} + \sqrt{y}}$

Q3 a) Test the consistency of [6]

$$2x - y - z = 2, \quad x + 2y + z = 2, \quad 4x - 7y - 5z = 2$$

And Solve if consistent.

b) Examine the function for its extreme values [6]

$$f(x, y) = y^2 + 4xy + 3x^2 + x^3$$

c) If $\sin(\theta + i\varphi) = e^{i\alpha}$ then Prove $\cos^4 \theta = \sin^2 \alpha = \sinh^4 \varphi$ [8]Q4 a) If $x = u \cos v$, $y = u \sin v$ then [6]

$$\text{Prove } \frac{\partial(u,v)}{\partial(x,y)} \cdot \frac{\partial(x,y)}{\partial(u,v)} = 1$$

b) If $\log(x + iy) = e^p(\cos q + i \sin q)$ then [6]
prove that $y = x \tan(\tan q \cdot \log \sqrt{x^2 + y^2})$

c) Solve by Gauss Elimination method [8]

$$2x + 3y + 4z = 11, \quad x + 5y + 7z = 1, \quad 3x + 11y + 13z = 25$$

- Q5 a) Prove $\cos^6 \theta + \sin^6 \theta = \frac{1}{8}[3 \cos 4\theta + 5]$ [6]
- b) Evaluate $\lim_{x \rightarrow 0} \left[\frac{1}{x^2} - \cot^2 x \right]$ [6]
- c) If $y = \cos(m \sin^{-1} x)$ then [8]
 prove that $(1 - x^2)y_{n+2} - (2n + 1)x y_{n+1} + (m^2 - n^2)y_n = 0$

- Q6 a) Check if the following vectors [6]

$$X_1 = [1, 0, 2, 1], X_2 = [3, 1, 2, 1], X_3 = [4, 6, 2, -4],$$

$X_4 = [-6, 0, -3, -4]$ are linear dependent hence find the relation between them if any.

- b) If $f(xy^2, z - 2x) = 0$ then [6]

$$\text{prove that } 2x \frac{\partial z}{\partial x} - y \frac{\partial z}{\partial y} = 4x$$

- c) Fit a second degree parabola $y = ax^2 + bx + c$ to the following data [8]

x	1	2	3	4	5	6	7	8	9
y	2	6	7	8	10	11	11	10	9